PB-245 168

LAKE NAOMI, MONROE COUNTY, PENNSYLVANIA

Pacific Northwest Environmental
Research Laboratory

Prepared for:

National Environmental Research Center

June 1975

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE

PB 245 168

U.S. ENVIRONMENTAL PROTECTION AGENCY
NATIONAL EUTROPHICATION SURVEY

WORKING PAPER SERIES

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
US Department of Commerce
Springfield, VA. 22151

PACIFIC NORTHWEST ENVIRONMENTAL RESEARCH LABORATORY

An Associate Laboratory of the

NATIONAL ENVIRONMENTAL RESEARCH CENTER - CORVALLIS, OREGON

and

NATIONAL ENVIRONMENTAL RESEARCH CENTER - LAS VEGAS, NEVADA

7-032

TECHNICAL REPORT DATA
(Please read Instructions on the reverse before completing)

1. REPORT NO.	2	3. RECIPIENT'S ACCESSION NO.
4. TITLE AND SUBTITLE Report on Lake Naomi, Monroe County, Pennsylvania		5. REPORT DATE Prepared June 1975 6. PERFORMING ORGANIZATION CODE
7. AUTHOR(S) Anonymous (National Eutrophication Survey Staff)		8. PERFORMING ORGANIZATION REPORT NO. Working Paper # 421
9. PERFORMING ORGANIZATION NAME AND ADDRESS National Eutrophication Survey U.S. Environmental Protection Agency 200 S. W. 35th St. Corvallis, Oregon 97330		10. PROGRAM ELEMENT NO. 11. CONTRACT/GRANT NO.
12. SPONSORING AGENCY NAME AND ADDRESS Office of Research and Development U.S. Environmental Protection Agency 401 M Street, S. W. Washington, DC 20460		13. TYPE OF REPORT AND PERIOD COVERED Final - 1973 14. SPONSORING AGENCY CODE

15. SUPPLEMENTARY NOTES
Prepared in cooperation with the U.S. Environmental Protection Agency, National Environmental Research Center, Las Vegas, Nevada 89114

16. ABSTRACT

Annual total phosphorus and total nitrogen loadings to the lake were estimated and subdivided according to either point or non-point source origin. An assessment of the lake's trophic condition and limiting nutrient is also provided. All data collected by the U.S.E.P.A. National Eutrophication Survey during the one year study of the lake and its tributaries are included in the report.

17. KEY WORDS AND DOCUMENT ANALYSIS

a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
eutrophication lakes nutrients phosphorus nitrogen loading		05B 05C

PRICES SUBJECT TO CHANGE

18. DISTRIBUTION STATEMENT	19. SECURITY CLASS (This Report)	21. NO. OF PAGES
Release unlimited		31
10	20. SECURITY CLASS (This page)	22. PRICE 3.75 · 2.25

EPA Form 2220-1 (9-73)

276054

REPORT
ON
LAKE NAOMI
MONROE COUNTY
PENNSYLVANIA
EPA REGION III
WORKING PAPER No. 421

WITH THE COOPERATION OF THE
PENNSYLVANIA DEPARTMENT OF ENVIRONMENTAL RESOURCES
AND THE
PENNSYLVANIA NATIONAL GUARD

JUNE, 1975

CONTENTS

	Page
Foreword	ii
List of Pennsylvania Study Lakes	iv
Lake and Drainage Area Map	v
Sections	
I. Conclusions	1
II. Lake and Drainage Basin Characteristics	3
III. Lake Water Quality Summary	4
IV. Nutrient Loadings	8
V. Literature Reviewed	13
VI. Appendices	14

FOREWORD

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nationwide threat of accelerated eutrophication to fresh water lakes and reservoirs.

OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.

b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.

c. With such a transformation, an assessment of the potential for eutrophication control can be made.

LAKE ANALYSIS

In this report, the first stage of evaluation of lake and watershed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

ACKNOWLEDGMENT

The staff of the National Eutrophication Survey (Office of Research & Development, U. S. Environmental Protection Agency) expresses sincere appreciation to the Pennsylvania Department of Environmental Resources for professional involvement and to the Pennsylvania National Guard for conducting the tributary sampling phase of the Survey.

Walter A. Lyon, Director of the Bureau of Water Quality Management, Richard M. Boardman, Chief of the Division of Water Quality, and James T. Ulanoski, Aquatic Biologist of the Division of Water Quality, provided invaluable lake documentation and counsel during the Survey, reviewed the preliminary reports, and provided critiques most useful in the preparation of this Working Paper series.

Major General Harry J. Mier, Jr., the Adjutant General of Pennsylvania, and Project Officer Major Ronald E. Wickard, who directed the volunteer efforts of the Pennsylvania National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.

NATIONAL EUTROPHICATION SURVEY

STUDY LAKES

STATE OF PENNSYLVANIA

LAKE NAME	COUNTY
Allegheny Reservoir	McKean, Warren, PA; Cattarugus, NY
Beaver Run Reservoir	Westmoreland
Beltzville	Carbon
Blanchard Reservoir	Centre
Canadohta	Crawford
Conneaut	Crawford
Conewago (Pinchot)	York
Greenlane	Montgomery
Harveys	Luzerne
Indian	Somerset
Naomi	Monroe
Ontelaunee	Berks
Pocono	Monroe
Pymatuning Reservoir	Crawford, PA; Ashtabula, OH
Shenango River Reservoir	Mercer
Stillwater	Monroe
Wallenpaupack	Pike, Wayne


LAKE NAOMI
Tributary Sampling Site
Lake Sampling Site
0 1 2Km.
0 1Mi.
Scale
41 08'
Dotters Run
Stillwater Lake
Upper Tunkhannock Creek
dam
24A2
28A1
NAOMI
LAKE
X2401
Upper Tunkhannock Creek
24A1 outlet
dam
Pocono Pines
Pocono Lake
Tamaqua Lake
Beaver Creek
41 06'
Penn.
Map Location
Lost Lakes
75 28'
75 26'

LAKE NAOMI*

STORET NO. 4224

I. CONCLUSIONS

A. Trophic Condition:

Survey data indicate that Lake Naomi is mesotrophic. It ranked first in overall trophic quality when the 17 Pennsylvania lakes sampled in 1973 were compared using a combination of six lake parameters**. Three of the lakes had less median total phosphorus, one had less and three had the same median dissolved phosphorus, two had less median inorganic nitrogen, five had less mean chlorophyll a, and nine had greater mean Secchi disc transparency.

Survey limnologists noted sparse rooted aquatic vegetation in July, 1973.

B. Rate-Limiting Nutrient:

The algal assay results indicate that Lake Naomi was limited by phosphorus at the time the sample was taken (04/17/73). The lake data indicate phosphorus limitation at all sampling times.

C. Nutrient Controllability:

1. Point sources--The estimated phosphorus load from point sources amounted to 8.0% of the total load. Of this estimated load, 5.4% was contributed by the Radio Church of God, and 2.7% was from shoreline septic tanks.

---

* Table of metric conversions--Appendix A.
** See Appendix B.

While the present phosphorus loading rate of 0.28 $g/m^2/yr$ is less than that proposed by Vollenweider (Vollenweider and Dillon, 1974) as an oligotrophic rate (see page 12), every effort should be made to reduce all phosphorus inputs to the lowest practicable level to protect the existing high quality of Lake Naomi.

2. Non-point sources--The phosphorus load from non-point sources, including precipitation, was 92.0% of the total load reaching the lake during the sampling year. Most of this load was contributed by Upper Tunkhannock Creek (63.4%). The ungaged drainage area was estimated to have contributed 22.3%.

The non-point phosphorus export of Upper Tunkhannock Creek (see page 12) is quite comparable to the exports of the two tributaries of nearby Stillwater Lake* (mean of 10 $kg/km^2/yr$; range of 8 to 12) and the three tributaries of downstream Pocono Lake* (mean of 10 $kg/km^2/yr$; range of 5 to 14).

* Respectively, Working Paper No. 427 and No. 424.

II. LAKE AND DRAINAGE BASIN CHARACTERISTICS

A. Lake Morphometry†:

1. Surface area: 2.02 kilometers^2.
2. Mean depth: 0.9 meters.
3. Maximum depth: 4.9 meters.
4. Volume: 1.818 x 10^6 m^3.
5. Mean hydraulic retention time: 21 days.

B. Tributary and Outlet:
(See Appendix C for flow data)

1. Tributaries -

Name	Drainage area (km^2)*	Mean flow (m^3/sec)*
Upper Tunkhannock Creek (A-2)	35.0	0.7
Minor tributaries & immediate drainage -	12.5	0.3
Total	47.5	1.0

2. Outlet -

Upper Tunkhannock Creek (A-1)	49.5**	1.0

C. Precipitation***:

1. Year of sampling: 149.0 centimeters.
2. Mean annual: 120.3 centimeters.

---

† Ulanoski, 1975.
* For limits of accuracy, see Working Paper No. 175, "...Survey Methods, 1973-1976".
** Includes area of lake.
*** See Working Paper No. 175.

III. LAKE WATER QUALITY SUMMARY

Lake Naomi was sampled three times during the open-water season of 1973 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from one or more depths at a single station on the lake (see map, page v). During each visit, a single depth-integrated (near bottom to surface) sample was collected for phytoplankton identification and enumeration; and a similar sample was collected for chlorophyll a analysis. During the first visit, an 18.9-liter depth-integrated sample was collected for algal assays. The maximum depth sampled was 1.2 meters.

The lake sampling results are presented in full in Appendix D and are summarized in the following table.

A. SUMMARY OF PHYSICAL AND CHEMICAL CHARACTERISTICS FOR LAKE NAOMI
STORET CODE 4224

PARAMETER	1ST SAMPLING (6/17/73) 1 SITES RANGE	MEAN	MEDIAN	2ND SAMPLING (7/23/73) 1 SITES RANGE	MEAN	MEDIAN	3RD SAMPLING (10/ 3/73) 1 SITES RANGE	MEAN	MEDIAN
TEMP (C)	10.9 - 10.9	10.9	10.9	22.7 - 22.7	22.7	22.7	17.1 - 17.1	17.1	17.1
DISS OXY (MG/L)	10.6 - 10.6	10.6	10.6	7.0 - 7.0	7.0	7.0	8.2 - 8.2	8.2	8.2
CNDCTVY (MCROMO)	50. - 50.	50.	50.	32. - 32.	32.	32.	27. - 27.	27.	27.
PH (STAND UNITS)	7.5 - 7.6	7.5	7.5	5.5 - 5.5	5.5	5.5	6.0 - 6.0	6.0	6.0
TOT ALK (MG/L)	10. - 10.	10.	10.	10. - 10.	10.	10.	10. - 10.	10.	10.
TOT P (MG/L)	0.011 - 0.011	0.011	0.011	0.020 - 0.020	0.020	0.020	0.017 - 0.017	0.017	0.017
ORTHO P (MG/L)	0.006 - 0.006	0.006	0.006	0.005 - 0.005	0.005	0.005	0.004 - 0.004	0.004	0.004
NO2+NO3 (MG/L)	0.100 - 0.100	0.100	0.100	0.090 - 0.090	0.090	0.090	0.030 - 0.030	0.030	0.030
AMMONIA (MG/L)	0.030 - 0.040	0.035	0.035	0.100 - 0.100	0.100	0.100	0.050 - 0.050	0.050	0.050
KJEL N (MG/L)	0.200 - 0.300	0.250	0.250	0.500 - 0.500	0.500	0.500	0.400 - 0.400	0.400	0.400
INORG N (MG/L)	0.130 - 0.140	0.135	0.135	0.190 - 0.190	0.190	0.190	0.080 - 0.080	0.080	0.080
TOTAL N (MG/L)	0.300 - 0.400	0.350	0.350	0.590 - 0.590	0.590	0.590	0.430 - 0.430	0.430	0.430
CHLRPYL A (UG/L)	1.8 - 1.8	1.8	1.8	10.5 - 10.5	10.5	10.5	4.3 - 4.3	4.3	4.3
SECCHI (METERS)	1.5 - 1.5	1.5	1.5	1.3 - 1.3	1.3	1.3	1.5 - 1.5	1.5	1.5

B. Biological characteristics:

1. Phytoplankton* -

Sampling Date	Dominant Genera	Algal units per ml
07/23/73	1. Synedra	1,784
	2. Melosira	758
	3. Glenodinium	558
	4. Asterionella	312
	5. Arthrodesmus	223
	Other genera	447
	Total	4,082
10/03/73	1. Flagellates	1,080
	2. Peridinium	163
	3. Cryptomonas	131
	4. Chlorophyta cells	65
	5. Asterionella	33
	Other genera	99
	Total	1,571

2. Chlorophyll a -

Sampling Date	Station Number	Chlorophyll a (µg/l)
04/17/73	01	1.8
07/23/73	01	10.5
10/03/73	01	4.3

C. Limiting Nutrient Study:

1. Autoclaved, filtered, and nutrient spiked -

Spike (mg/l)	Ortho P Conc. (mg/l)	Inorganic N Conc. (mg/l)	Maximum yield (mg/l-dry wt.)	N/P Ratio
Control	<0.005	0.126	0.1	25/1
0.050 P	0.055	0.126	1.2	2/1
0.050 P + 1.0 N	0.055	1.126	7.6	20/1
1.0 N	<0.005	1.126	0.1	225/1

* The April sample was lost in shipment.

2. Discussion -

The control yield of the assay alga, Selenastrum capricornutum, indicates that the potential primary productivity of Lake Naomi was low at the time the assay sample was collected. The relatively small but significant increase in yield when orthophosphorus was added and the lack of change in yield when only nitrogen was added indicate phosphorus limitation.

The relatively small increase in yield resulting from the orthophosphorus spike was due to the shift to nitrogen limitation when the excess phosphorus was added (note the change in the N/P ratios). The control sample would have become nitrogen limited with the addition of only about 0.010 mg/l of orthophosphorus, so 4/5ths of the 0.050 mg/l spike could not be utilized by the assay algae. Hence the small growth response to the phosphorus addition.

The lake data indicate phosphorus limitation at all sampling times (the mean N/P ratios were 20/1 or greater, and phosphorus limitation would be expected).

IV. NUTRIENT LOADINGS
(See Appendix E for data)

For the determination of nutrient loadings, the Pennsylvania National Guard collected monthly near-surface grab samples from each of the tributary sites indicated on the map (page v), except for the high runoff months of February and March when two samples were collected. Sampling was begun in May, 1973, and was completed in April, 1974.

Through an interagency agreement, stream flow estimates for the year of sampling and a "normalized" or average year were provided by the Pennsylvania District Office of the U.S. Geological Survey for the tributary sites nearest the lake.

In this report, nutrient loads for sampled tributaries were determined by using a modification of a U.S. Geological Survey computer program for calculating stream loadings*. Nutrient loads shown are those measured minus point-source loads, if any.

Nutrient loads for unsampled "minor tributaries and immediate drainage" ("ZZ" of U.S.G.S.) were estimated using the means of the nutrient loads, in $kg/km^2/year$, at station A-2 and multiplying the means by the ZZ area in km^2.

The Radio Church of God did not participate in the Survey, and nutrient loads were estimated at 1.134 kg P and 3.401 kg N/capita/year.

---

* See Working Paper No. 175.

A. Waste Sources:

1. Known muncipal* -

Name	Pop. Served	Treatment	Mean Flow (m^3/d)	Receiving Water
Radio Church of God	300**	lagoon	?	Lake Naomi

2. Known industrial - None

* Ulanoski, 1975.
** Population is seasonal with a maximum of 300 persons for approximately one month.

B. Annual Total Phosphorus Loading - Average Year:

1. Inputs -

Source	kg P/yr	% of total
a. Tributaries (non-point load) -		
Upper Tunkhannock Creek (A-2)	355	63.4
b. Minor tributaries & immediate drainage (non-point load) -	125	22.3
c. Known STP's -		
Radio Church of God	30	5.4
d. Septic tanks* -	15	2.7
e. Known industrial - None	-	-
f. Direct precipitation** -	35	6.2
Total	560	100.0

2. Outputs -

Lake outlet - Upper Tuckhannock Creek (A-1)	540

3. Net annual P accumulation - 20 kg.

* Estimate based on 60 shoreline dwellings; see Working Paper No. 175.
** See Working Paper No. 175.

C. Annual Total Nitrogen Loading - Average Year:

1. Inputs -

Source	kg N/yr	% of total
a. Tributaries (non-point load) -		
Upper Tunkhannock Creek (A-2)	13,820	63.8
b. Minor tributaries & immediate drainage (non-point load) -	4,940	22.8
c. Known STP's -		
Radio Church of God	85	0.4
d. Septic tanks* -	640	3.0
e. Known industrial - None	-	-
f. Direct precipitation** -	2,180	10.0
Total	21,665	100.0

2. Outputs -

Lake outlet - Upper Tunkhannock Creek (A-1)	17,695

3. Net annual N accumulation - 3,970 kg.

* Estimate based on 60 shoreline dwellings; see Working Paper No. 175.
** See Working Paper No. 175.

D. Mean Annual Non-point Nutrient Export by Subdrainage Area:

Tributary	kg $P/km^2/yr$	kg $N/km^2/yr$
Upper Tunkhannock Creek (A-2)	10	395

E. Yearly Loading Rates:

In the following table, the existing phosphorus loading rates are compared to those proposed by Vollenweider (Vollenweider and Dillon, 1974). Essentially, his "dangerous" rate is the rate at wnich the receiving water would become eutrophic or remain eutrophic; his "permissible" rate is that which would result in the receiving water remaining oligotrophic or becoming oligotrophic if morphometry permitted. A mesotrophic rate would be considered one between "dangerous" and "permissible".

Note that Vollenweider's model may not be applicable to water bodies with short hydraulic retention times.

	Total Phosphorus		Total Nitrogen	
	Total	Accumulated	Total	Accumulated
grams/m^2/yr	0.28	0.01	10.7	2.0

Vollenweider loading rates for phosphorus ($g/m^2/yr$) based on mean depth and mean hydraulic retention time of Lake Naomi:

"Dangerous" (eutrophic rate)	0.74
"Permissible" (oligotrophic rate)	0.37

V. LITERATURE REVIEWED

Ulanoski, James T., 1975. Personal communication (point sources at Lake Naomi). PA Dept. of Env. Resources, Harrisburg.

Vollenweider, R. A., and P. J. Dillon, 1974. The application of the phosphorus loading concept to eutrophication research. Natl. Res. Council of Canada Publ. No. 13690, Canada Centre for Inland Waters, Burlington, Ontario.

CONVERSION FACTORS

Hectares x 2.471 = acres

Kilometers x 0.6214 = miles

Meters x 3.281 = feet

Cubic meters x 8.107×10^{-4} = acre/feet

Square kilometers x 0.3861 = square miles

Cubic meters/sec x 35.315 = cubic feet/sec

Centimeters x 0.3937 = inches

Kilograms x 2.205 = pounds

Kilograms/square kilometer x 5.711 = lbs/square mile

LAKES RANKED BY INDEX NOS.

RANK	LAKE CODE	LAKE NAME	INDEX NO
1	4224	LAKE NAOMI	445
2	4220	BELTZVILLE DAM	423
3	4222	HARVEY'S LAKE	413
4	4228	STILLWATER LAKE	401
5	4227	POCONO LAKE	389
6	4223	INDIAN LAKE	388
7	3641	ALLEGHENY RESERVOIR	385
8	4229	LAKE WALLENPAUPACK	371
9	4221	CANADOHTA LAKE	369
10	4219	BEAVER RUN RESERVOIR	360
11	4204	CONNEAUT LAKE	307
12	4226	PINCHOT LAKE	256
13	4213	PYMATUNING RESERVOIR	206
14	4216	SHENANGO RIVER RESERVOIR	157
15	4225	ONTELAUNEE DAM	101
16	4201	BLANCHARD RESERVOIR	85
17	4207	GREENLANE DAM	53

PERCENT OF LAKES WITH HIGHER VALUES (NUMBER OF LAKES WITH HIGHER VALUES)

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500-MEAN SEC	MEAN CHLORA	15-MIN DO	MEDIAN DISS P	INDEX NO
3641	ALLEGHENY RESERVOIR	56 (9)	38 (6)	63 (10)	100 (16)	69 (11)	50 (8)	385
4201	BLANCHARD RESERVOIR	13 (2)	13 (2)	25 (4)	31 (5)	3 (0)	0 (0)	85
4204	CONNEAUT LAKE	44 (7)	63 (10)	69 (11)	56 (9)	34 (5)	41 (6)	307
4207	GREENLANE DAM	6 (1)	6 (1)	19 (3)	13 (2)	3 (0)	6 (1)	53
4213	PYMATUNING RESERVOIR	0 (0)	72 (11)	6 (1)	0 (0)	100 (16)	28 (4)	206
4216	SHENANGO RIVER RESERVOIR	19 (3)	44 (7)	13 (2)	6 (1)	47 (7)	28 (4)	157
4219	BEAVER RUN RESERVOIR	94 (15)	19 (3)	88 (14)	81 (13)	19 (2)	59 (8)	360
4220	BELTZVILLE DAM	88 (14)	25 (4)	94 (15)	94 (15)	34 (5)	88 (13)	423
4221	CANADOHTA LAKE	50 (8)	97 (15)	56 (9)	19 (3)	59 (9)	88 (13)	369
4222	HARVEY'S LAKE	63 (10)	81 (13)	100 (16)	63 (10)	47 (7)	59 (8)	413
4223	INDIAN LAKE	100 (16)	31 (5)	75 (12)	75 (12)	19 (2)	88 (13)	388
4224	LAKE NAOMI	81 (13)	88 (14)	44 (7)	69 (11)	88 (14)	75 (12)	445
4225	ONTELAUNEE DAM	25 (4)	0 (0)	0 (0)	44 (7)	19 (2)	13 (2)	101
4226	PINCHOT LAKE	31 (5)	56 (9)	31 (5)	38 (6)	81 (13)	19 (3)	256
4227	POCONO LAKE	38 (6)	97 (15)	50 (8)	88 (14)	75 (12)	41 (6)	389
4228	STILLWATER LAKE	72 (11)	72 (11)	38 (6)	25 (4)	94 (15)	100 (16)	401
4229	LAKE WALLENPAUPACK	72 (11)	50 (8)	81 (13)	50 (8)	59 (9)	59 (8)	371

LAKE DATA TO BE USED IN RANKINGS

LAKE CODE	LAKE NAME	MEDIAN TOTAL P	MEDIAN INORG N	500-MEAN SEC	MEAN CHLORA	15-MIN DO	MEDIAN DISS P
3641	ALLEGHENY RESERVOIR	0.016	0.380	414.250	3.700	13.800	0.006
4201	BLANCHARD RESERVOIR	0.064	1.300	453.143	15.187	14.900	0.046
4204	CONNEAUT LAKE	0.023	0.185	402.000	7.567	14.600	0.007
4207	GREENLANE DAM	0.066	1.475	460.222	24.011	14.900	0.020
4213	PYMATUNING RESERVOIR	0.070	0.180	467.750	56.333	7.700	0.008
4216	SHENANGO RIVER RESERVOIR	0.058	0.340	463.555	26.800	14.500	0.008
4219	BEAVER RUN RESERVOIR	0.009	0.835	384.833	5.183	14.800	0.006
4220	BELTZVILLE DAM	0.010	0.815	362.444	4.856	14.600	0.005
4221	CANADOHTA LAKE	0.020	0.130	436.000	19.167	14.100	0.005
4222	HARVEY'S LAKE	0.015	0.160	338.000	5.967	14.500	0.006
4223	INDIAN LAKE	0.008	0.520	400.222	5.211	14.800	0.005
4224	LAKE NAOMI	0.014	0.135	443.333	5.533	8.000	0.005
4225	ONTELAUNEE DAM	0.040	2.150	470.667	11.783	14.800	0.011
4226	PINCHOT LAKE	0.027	0.245	453.000	13.950	11.500	0.008
4227	POCONO LAKE	0.024	0.130	438.800	4.980	13.200	0.007
4228	STILLWATER LAKE	0.015	0.180	449.000	18.233	7.900	0.004
4229	LAKE WALLENPAUPACK	0.015	0.250	394.583	9.617	14.100	0.006

TRIBUTARY FLOW INFORMATION FOR PENNSYLVANIA 1/27/75

LAKE CODE 4224 NAUMI LAKE

TOTAL DRAINAGE AREA OF LAKE(SQ KM) 49.5

TRIBUTARY	SUB-DRAINAGE AREA(SQ KM)	NORMALIZED FLOWS(CMS) JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	MEAN
4224A1	44.5	0.98	3.08	1.67	1.34	1.36	0.82	0.65	0.59	0.34	0.54	0.93	1.10	0.99
4224A2	35.0	0.68	0.68	1.05	1.16	0.84	0.54	0.51	0.45	0.31	0.42	0.68	0.76	0.68
4224Z2	14.5	0.24	0.28	0.42	0.42	0.48	0.23	0.14	0.14	0.03	0.11	0.25	0.34	0.31

SUMMARY

TOTAL DRAINAGE AREA OF LAKE = 49.5
SUM OF SUB-DRAINAGE AREAS = 49.5
TOTAL FLOW IN = 11.92
TOTAL FLOW OUT = 11.92

MEAN MONTHLY FLOWS AND DAILY FLOWS(CMS)

TRIBUTARY	MONTH	YEAR	MEAN FLOW	DAY	FLOW	DAY	FLOW	DAY	FLOW
4224A1	5	73	1.48	17	6.75				
	6	73	2.55	23	9.79				
	7	73	1.13	14	6.96				
	8	73	0.71	4	1.70				
	9	73	0.42	8	0.65				
	10	73	0.99	13	0.31				
	11	73	0.83	10	0.91				
	12	73	3.17	4	6.08				
	1	74	1.61	12	1.98				
	2	74	1.13	9	0.82				
	3	74	1.34	10	1.93				
	4	74	1.65	13	1.90				
4224A2	5	73	1.33	17	1.75				
	6	73	1.54	23	0.82				
	7	73	0.85	14	0.74				
	8	73	0.57	4	1.16				
	9	73	0.31	8	0.69				
	10	73	0.96	13	0.28				
	11	73	0.81	10	0.42				
	12	73	1.45	8	1.81				
	1	74	1.12	12	0.79				
	2	74	0.82	9	0.81				
	3	74	1.12	15	1.15				
	4	74	1.31	13	1.61				
4224Z2	5	73	0.54	17	0.95				
	6	73	0.71	23	0.22				
	7	73	0.63	14	0.21				
	8	73	0.17	4	0.49				
	9	73	0.13	8	0.02				
	10	73	0.19	13	0.07				
	11	73	0.15	10	0.31				
	12	73	0.49	8	0.79				
	1	74	0.35	12	0.51				
	2	74	0.34	9	0.15				
	3	74	0.43	15	0.45				
	4	74	0.55	13	0.39				

APPENDIX D

PHYSICAL and CHEMICAL DATA

STORET RETRIEVAL DATE 75/01/27

422401
41 06 35.0 075 27 36.0
LAKE NAOMI
42089 PENNSYLVANIA

11EPALES 2111202
3 0008 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00010 WATER TEMP CENT	00300 DO MG/L	00077 TRANSP SECCHI INCHES	00094 CNDUCTVY FIELD MICROMHO	00400 PH SU	00410 T ALK CACO3 MG/L	00610 NH3-N TOTAL MG/L	00625 TOT KJEL N MG/L	00630 NO2&NO3 N-TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P
73/04/17	10 30	0000	10.9		60	50K	7.60	10K	0.040	0.300	0.100	0.006
	10 30	0004	10.9	10.6		50K	7.50	10K	0.030	0.200	0.100	0.006
73/07/23	10 30	0000	22.7	7.0	50	32	5.50	10K	0.100	0.500	0.090	0.005
73/10/03	16 00	0000	17.1	8.2	60	27	6.00	10K	0.050	0.400	0.030	0.004

20

DATE FROM TO	TIME OF DAY	DEPTH FEET	00665 PHOS-TOT MG/L P	32217 CHLRPHYL A UG/L
73/04/17	10 30	0000	0.011	1.8
	10 30	0004	0.011	
73/07/23	10 30	0000	0.020	10.5
73/10/03	16 00	0000	0.017	4.3

K VALUE KNOWN TO BE
LESS THAN INDICATED

APPENDIX E

TRIBUTARY DATA

STORET RETRIEVAL DATE 75/02/03

4224A1
41 06 30.0 075 28 30.0
UPPER TUNKHANNOCK CR
42111 7.5 POCONO PINES
U/NAOMI LAKE
HWY 423 BRDG IN TOWN OF PONOCO LAKE
11EPALES 2111204
4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 NO2&NO3 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/05/14	15 37		0.058	0.310	0.005K	0.005K	0.015
73/06/23	10 25		0.024	0.440	0.029	0.005K	0.025
73/07/14	09 20		0.012	0.750	0.078	0.010	0.020
73/08/05	10 15		0.021	0.320	0.022	0.005K	0.025
73/09/09	10 10		0.019	1.600	0.080	0.005K	0.025
73/10/07	11 10		0.013	0.350	0.020	0.006	0.012
73/11/11	14 00		0.024	1.050	0.052		0.020
73/12/09	12 40		0.092	0.300	0.020	0.005K	0.020
74/01/06	10 50		0.136	0.100	0.024	0.005K	0.010
74/02/04	14 45		0.184	0.400	0.035	0.005K	0.005
74/02/23	14 45		0.192	0.300	0.025	0.010	0.020
74/03/16	10 10		0.152	0.500	0.045	0.005K	0.015
74/03/30	14 00		0.144	0.250	0.015	0.005	0.012
74/04/13	10 20		0.092	0.100	0.010	0.005K	0.010

22

K VALUE KNOWN TO BE
LESS THAN INDICATED

STORET RETRIEVAL DATE 75/02/03

4224A2
41 07 02.0 075 25 31.0
UPPER TUNKHANNOCK CR
42 7.5 POCONO PINES
1/NAOMI LAKE
RD BRDG JUST BELO STILLWATER LAKE OUTLET
11EPALES 2111204
4 0000 FEET DEPTH

DATE FROM TO	TIME OF DAY	DEPTH FEET	00630 NO2&NO3 N-TOTAL MG/L	00625 TOT KJEL N MG/L	00610 NH3-N TOTAL MG/L	00671 PHOS-DIS ORTHO MG/L P	00665 PHOS-TOT MG/L P
73/05/19	15 30		0.056	0.340	0.005K	0.005K	0.015
73/06/23	10 05		0.011	0.400	0.020	0.005K	0.020
73/07/14	10 00		0.010K	0.630	0.084	0.008	0.015
73/08/05	10 30		0.010K	0.240	0.018	0.005K	0.015
73/09/09	10 00		0.012	2.100	0.126	0.005K	0.015
73/10/07	10 47		0.010K	0.400	0.034	0.006	0.010
73/11/11	13 15		0.056	1.000	0.052	0.016	0.020
73/12/04	11 15		0.104	0.300	0.024	0.008	0.015
74/01/06	10 06		0.136	0.400	0.040	0.005K	0.010
74/02/09	14 27		0.168	0.600	0.055	0.005K	0.010
74/02/23	14 27		0.152	0.200	0.010	0.010	0.025
74/03/16	09 40		0.168	0.800	0.055	0.005	0.025
74/03/30	12 00		0.152	0.300	0.035	0.005	0.010
74/04/13	11 40		0.116	0.200	0.015	0.005K	0.015

K VALUE KNOWN TO BE
LESS THAN INDICATED